

**S8.P24****Role of DsrC in the mechanism of sulfite reduction by the dissimilatory sulfite reductase DsrAB**

André A. Santos<sup>a</sup>, Sofia S. Venceslau<sup>a</sup>, Fabian Grein<sup>b</sup>, Inês Cardoso Pereira<sup>a</sup>  
<sup>a</sup>ITQB-UNL, Portugal

<sup>b</sup>University of Bonn, Germany

E-mail: [andresantos@itqb.unl.pt](mailto:andresantos@itqb.unl.pt)

The dissimilatory sulfite reductase (DsrAB) is one of the most important enzymes in the biogeochemical sulfur cycle [1]. It is present in all sulfate reducing microorganisms, in sulfite/thiosulfate/organosulfonate reducers and in sulfur-oxidizing bacteria. This siroheme-containing enzyme catalyzes the reduction of sulfite, but despite years of research, the mechanism and physiological products of sulfite reduction have not been clearly identified. It is not clear why in vitro DsrAB produces a mixture of products including thiosulfate and trithionate, while the closely-related assimilatory enzyme reduces sulfite directly to sulfide. More recently, the reduction of sulfite by DsrAB was proposed to involve also the small protein DsrC, which contains two conserved redoxactive cysteines in a flexible C-terminal arm [2]. The crystal structure of DsrAB in complex with DsrC showed that one of these Cys is located right next to the active site, pointing to the involvement of DsrC in the reduction mechanism [3]. Here, we report recent results from the investigation on the role of DsrC in the reduction of sulfite by DsrAB, using in vivo and in vitro experiments. Our results show that DsrC is directly involved in sulfite reduction and permits the identification of the mechanism and physiological product of this important reaction.

**References**

- [1] Grein F, Ramos AR, Venceslau SS, Pereira IAC (2013) Unifying concepts in anaerobic respiration: Insights from dissimilatory sulfur metabolism. *Biochim Biophys. Acta-Bioenergetics* 1827, 145–60.
- [2] Venceslau SS, Stockdreher Y, Dahl C, Pereira IAC (2014), The “bacterial heterodisulfide” DsrC is a key protein in dissimilatory sulfur metabolism, *Biochim. Biophys. Acta-Bioenergetics*, <http://dx.doi.org/10.1016/j.bbabi.2014.03.007>.
- [3] Oliveira TF, Vornheim, Matias PM, Venceslau SS, Pereira IAC, Archer M (2008) The crystal structure of a dissimilatory sulfite reductase bound to DsrC provides novel insights into the mechanism of sulfate reduction. *J. Biol. Chem.* 283: 34141.

doi:[10.1016/j.bbabi.2014.05.105](https://doi.org/10.1016/j.bbabi.2014.05.105)

**S8.P25****Dithiothreitol, redox reagent, is affecting proton-translocating ATPase activity of *Rhodobacter sphaeroides***

Harutyun Sargsyan<sup>a</sup>, Lilit Gabrielyan<sup>a</sup>, Armen Trchounian<sup>b</sup>

<sup>a</sup>Department of Microbiology, Microbes and Plants Biotechnology, Faculty of Biology, Yerevan State University, Armenia

<sup>b</sup>Department of Microbiology, Plants & Microbes Biotechnology, Faculty of Biology, Yerevan State University, Armenia

E-mail: [hsargsyan@ysu.am](mailto:hsargsyan@ysu.am)

The H<sup>+</sup>-translocating FoF<sub>1</sub>-ATPase of purple bacteria, the main membrane-associated enzyme of bioenergetics relevance, can generate proton motive force under various conditions [1]. Its role in photo-fermentation and hydrogen production by these bacteria is suggested [2]. In the current work we have studied effects of DL-dithiothreitol (DTT), a redox reagent reducing disulfides, on N,N'-

dicyclohexylcarbodiimide (DCCD)-inhibited ATPase activity of purple bacteria *Rhodobacter sphaeroides* str. MDC6521 (from mineral spring waters in Armenian mountains) membrane vesicles for revealing the regulatory pathways of bacterial redox sensing. The membrane vesicles showed pronounced H<sup>+</sup>-translocating ATPase activity, which was inhibited on ~42% by 0.1 mM DCCD. After treatment of membrane vesicles with 0.5–2 mM DTT an additional increase (~1.6–1.8-folds) of DCCD-inhibited ATPase activity was observed. The increase of the ATPase activity by DTT under reducing conditions might be connected with the change of dithiol/disulfide status of this enzyme due to the fact that redox-sensitive thiols gain access to the ATPase. Therefore, the FoF<sub>1</sub>-ATPase might have an essential role in redox sensing of *R. sphaeroides*.

**References**

- [1] L. Hakobyan, L. Gabrielyan, A. Trchounian, Proton motive force in *Rhodobacter sphaeroides* under anaerobic conditions in the dark, *Curr. Microbiol.* 62 (2011) 415–419.
- [2] L. Hakobyan, L. Gabrielyan, A. Trchounian, Relationship of proton motive force and the FoF<sub>1</sub>-ATPase with bio-hydrogen production activity of *Rhodobacter sphaeroides*: effects of diphenylene iodonium, hydrogenase inhibitor, and its solvent dimethylsulphoxide, *J. Bioenerg. Biomembr.* 44 (2012) 495–502.

doi:[10.1016/j.bbabi.2014.05.106](https://doi.org/10.1016/j.bbabi.2014.05.106)

**S8.P26****Study of the respiratory arsenate reductase from *Halorhodospira halophila* definitively clarifies the evolutionary history of this versatile enzyme**

Barbara Schoepp-Cothenet<sup>a</sup>, Marielle Bauzan<sup>b</sup>, Anne-Lise Ducluzeau<sup>c</sup>, Fabien Pierrel<sup>d</sup>, Wolfgang Nitschke<sup>a</sup>

<sup>a</sup>Bioénergétique et Ingénierie des Protéines, UMR 7281, CNRS, Aix Marseille University, IMM FR3479, Marseille, France

<sup>b</sup>Institut de Microbiologie de la Méditerranée, FR3479, F-13402 Marseille Cedex 20, France

<sup>c</sup>Beadle Center, University of Nebraska-Lincoln, 1901 Vine Street, Lincoln, NE 68588-0660, USA

<sup>d</sup>Chimie et Biologie des Métaux, UMR 5249, CEA, CNRS, JF University, Grenoble, France

E-mail: [schoepp@imm.cnrs.fr](mailto:schoepp@imm.cnrs.fr)

The three presently known enzymes responsible for arsenic-using bioenergetic processes are arsenite oxidase (Aio), arsenate reductase (Arr) and alternative arsenite oxidase (Arx), all of which are molybdoenzymes from the vast group referred to as the Mo/W-bisPGD enzyme superfamily. Since arsenite is present in substantial amounts in hydrothermal environments (frequently considered as vestiges of primordial biochemistry), arsenite-based bioenergetics has early on been predicted to be ancient. Conflicting scenarios, however, have been put forward proposing either Arr/Arx or Aio as operating in the ancestral metabolism. Phylogenetic data argue in favour of Aio whereas biochemical and physiological data led several authors to propose the Arr/Arx enzyme as the most ancient anaerobic arsenite metabolising enzyme. Here we combine phylogenetic approaches with physiological and biochemical experiments to demonstrate that the Arr/Arx enzyme could not have been functional in the Archaea. We show that Arr reacts with menaquinones to reduce arsenate whereas Arx reacts with ubiquinone to oxidise arsenite, in line with thermodynamic considerations. The phylogeny of the quinone biosynthesis pathway, however, clearly indicates that the ubiquinone pathway is recent. An updated phylogeny of Arr/Arx furthermore indicates a recent emergence of this enzyme. We